
Nanda Nursing Diagnosis For Kidney Failure

*Nanda Nursing
Diagnosis For Kidney
Failure*

*Downloaded from
app.ardentx.com by
Vazquez & Finley*

UNDERSTANDING NANDA NURSING DIAGNOSIS FOR KIDNEY FAILURE: A COMPREHENSIVE GUIDE FOR CARE PLANNING

Kidney failure, also known as end-stage renal disease (ESRD), represents a serious and life-altering condition that affects millions of people worldwide. When the kidneys lose their ability to filter waste products, maintain fluid balance, and regulate electrolytes, the entire body suffers. For nurses, managing these complex patients requires more than just clinical skills—it demands a structured, evidence-based approach to care. This is precisely where a NANDA nursing diagnosis for kidney failure becomes an indispensable tool. By using standardized nursing diagnoses, healthcare professionals can identify patient problems accurately, prioritize interventions, and improve outcomes. In this article, we will explore the most relevant NANDA nursing diagnoses for kidney failure, examine their defining characteristics, and discuss how to implement them in real-world clinical settings.

WHAT IS A NANDA NURSING DIAGNOSIS?

Before diving into specific diagnoses, it is essential to understand what a NANDA nursing diagnosis actually is. NANDA

International, formerly the North American Nursing Diagnosis Association, developed a standardized classification system that allows nurses to identify and label patient problems in a consistent, evidence-based manner. Unlike medical diagnoses, which focus on disease processes, nursing diagnoses address the patient's response to those processes. For example, while a physician might diagnose chronic kidney disease, a nurse would use a NANDA nursing diagnosis such as "Excess Fluid Volume" or "Risk for Electrolyte Imbalance."

These diagnoses form the foundation of the nursing care plan. They guide assessment, intervention, and evaluation, ensuring that care is holistic and patient-centered. When applied to kidney failure, NANDA diagnoses help nurses address the physiological, psychological, and social challenges that patients face throughout their illness.

WHY KIDNEY FAILURE REQUIRES SPECIALIZED NURSING DIAGNOSES

Kidney failure is not a single event but a progressive condition with far-reaching consequences. As renal function declines, patients experience a cascade of complications: fluid overload, electrolyte disturbances, anemia, bone disease, cardiovascular strain, and cognitive changes. Each of these complications presents a unique set of

nursing challenges. Without a clear diagnostic framework, it is easy to miss subtle changes or prioritize the wrong interventions. A well-chosen NANDA nursing diagnosis for kidney failure keeps the care plan focused, measurable, and adaptable as the patient's condition evolves.

Moreover, kidney failure often requires complex treatments such as hemodialysis, peritoneal dialysis, or kidney transplantation. Each treatment modality introduces new risks and care needs. For instance, a patient undergoing hemodialysis may face vascular access complications, while a peritoneal dialysis patient might be at higher risk for infection. The appropriate NANDA diagnoses capture these nuances and help nurses tailor their care accordingly.

COMMON NANDA NURSING DIAGNOSES FOR KIDNEY FAILURE

The following are some of the most frequently used NANDA nursing diagnoses for patients with kidney failure. Each diagnosis is accompanied by its defining characteristics, related factors, and suggestions for nursing interventions.

Excess Fluid Volume

One of the hallmark problems in kidney failure is the inability to excrete adequate amounts of fluid. This leads to fluid retention, which can manifest as edema, hypertension, pulmonary congestion, and weight gain. The NANDA nursing diagnosis "Excess Fluid Volume" is defined as increased isotonic fluid retention. It is one of the most common

and critical diagnoses in this patient population.

Defining characteristics: The nurse may observe peripheral edema, pitting edema in dependent areas, jugular vein distension, crackles in the lungs, dyspnea, orthopnea, rapid weight gain, and decreased urine output. Patients often report shortness of breath, especially when lying flat, and a feeling of fullness or tightness in the body.

Related factors: The primary cause is compromised renal function, but contributing factors may include excessive fluid intake, noncompliance with fluid restrictions, and the use of medications that promote fluid retention.

Nursing interventions: Accurate daily weights are essential—an increase of 1 to 2 kilograms in 24 hours suggests significant fluid retention. Strict intake and output monitoring, along with daily assessment for edema and lung sounds, should be standard. Nurses also play a key role in educating patients about fluid restrictions, typically 1 to 1.5 liters per day depending on residual renal function and dialysis adequacy. Diuretics may be prescribed, but their effectiveness diminishes as kidney function declines.

Risk for Electrolyte Imbalance

Electrolyte disturbances are nearly universal in kidney failure. The most dangerous imbalances include hyperkalemia, hyperphosphatemia, hypocalcemia, and metabolic acidosis. The NANDA nursing diagnosis "Risk for Electrolyte Imbalance" captures the

patient's susceptibility to these life-threatening complications.

Risk factors: Decreased glomerular filtration rate, use of potassium-sparing diuretics, metabolic acidosis, and dietary noncompliance are significant contributors. Patients on hemodialysis are particularly vulnerable between sessions when waste products accumulate.

Nursing assessments: Laboratory monitoring is crucial. Serum potassium, phosphate, calcium, and bicarbonate levels should be checked regularly, especially before and after dialysis. Electrocardiogram changes such as peaked T waves in hyperkalemia or prolonged QT intervals in hypocalcemia may be early warning signs. Nurses should also assess for muscle weakness, cardiac irregularities, and neurological changes.

Interventions: Dietary education is a cornerstone of care. Patients need to understand which foods are high in potassium, phosphorus, and sodium. For hyperkalemia, emergency interventions may include administration of calcium gluconate, insulin and glucose, or potassium-binding resins. In chronic management, phosphate binders taken with meals help control serum phosphate levels.

Impaired Urinary Elimination

Kidney failure often results in a drastic reduction in urine output, a condition known as oliguria or anuria. The NANDA diagnosis "Impaired Urinary Elimination" refers to a disturbance in urine excretion. In the context of kidney

failure, this diagnosis acknowledges both the physical changes and the patient's emotional response to losing a basic bodily function.

Defining characteristics: Patients may report decreased urine frequency, scanty amounts of urine, or complete absence of urine output. Some patients experience nocturia in earlier stages, only to progress to oliguria later. Discolored urine or the presence of sediment may also be noted.

Related factors: The root cause is the progressive loss of nephron function. Additionally, obstruction from kidney stones or tumors may contribute in some cases.

Nursing care: Accurate measurement of urine output is essential, and a catheter may be necessary for precise monitoring in critically ill patients. Nurses should also provide emotional support, as the loss of urinary function can be distressing. Education about the purpose of dialysis as a replacement for kidney function can help patients adjust to their new reality.

Fatigue

Fatigue is one of the most common and debilitating symptoms of kidney failure. The NANDA nursing diagnosis "Fatigue" is defined as an overwhelming, sustained sense of exhaustion and decreased capacity for physical and mental work. This is not simply tiredness that resolves with rest; it is a profound, persistent lack of energy that affects every aspect of daily life.

Defining characteristics: Patients may report constant tiredness, lack of energy, inability to maintain usual routines, weakness, and difficulty concentrating.

They may need to nap frequently and still feel unrefreshed. Emotional irritability and decreased libido are also common.

Related factors: Anemia due to decreased erythropoietin production is a major contributor. Other factors include accumulation of metabolic waste products, poor sleep quality, depression, malnutrition, and the energy demands of dialysis itself.

Nursing interventions: Addressing anemia with erythropoiesis-stimulating agents and iron supplementation is often the first step. Nurses should also assess sleep patterns and recommend sleep hygiene strategies. Energy conservation techniques, such as prioritizing activities and pacing oneself, can be helpful. Encouraging gentle exercise, such as walking or stretching, may paradoxically improve energy levels over time. Emotional support and screening for depression are also important.

Risk for Infection

Patients with kidney failure are at increased risk for infection due to multiple factors: altered immune function, frequent hospitalizations, invasive procedures, and the presence of vascular access devices. The NANDA diagnosis "Risk for Infection" captures this vulnerability.

Risk factors: For patients on hemodialysis, the arteriovenous fistula, graft, or central venous catheter is a common entry point for bacteria. Peritoneal dialysis patients face the risk of peritonitis. Additionally, uremia itself impairs neutrophil and lymphocyte function, making the entire body more susceptible.

Nursing assessments: Daily inspection of access sites for redness, swelling, warmth, or drainage is essential. Vital signs should be monitored for fever or hypothermia. Laboratory indicators such as an elevated white blood cell count or positive cultures confirm infection. Patients should be educated about signs of infection and when to seek medical attention.

Interventions: Strict adherence to sterile technique during catheter care and dialysis initiation is nonnegotiable. Hand hygiene, proper wound care, and timely administration of prophylactic antibiotics when indicated are critical. For peritoneal dialysis patients, education about proper exchange technique cannot be overemphasized.

Impaired Tissue Integrity

Patients with kidney failure often develop skin problems due to edema, uremic toxins, and poor circulation. Pruritus, or severe itching, is particularly distressing. The NANDA diagnosis "Impaired Tissue Integrity" addresses damage to mucous membranes or integumentary tissues.

Defining characteristics: The nurse may note dry, flaky skin, excoriations from scratching, rashes, bruising, or delayed wound healing. Patients report intense itching, often worse at night or after dialysis. Skin may appear pallid or hyperpigmented in some areas.

Related factors: Uremic toxins that accumulate in the blood are deposited in the skin, causing irritation. Dry skin results from decreased sweat gland function and fluid loss during dialysis.

Additionally, calcium-phosphate imbalances can lead to pruritus.

Nursing interventions: Emollients and moisturizers help combat dryness. Cool compresses and antihistamines may relieve itching. Nurses should educate patients to avoid hot water, harsh soaps, and excessive scratching. Keeping nails trimmed and wearing soft fabrics can reduce skin trauma. In severe cases, topical steroids or ultraviolet therapy may be considered.

Ineffective Therapeutic Regimen Management

Kidney failure requires a complex and demanding self-care regimen: dietary restrictions, fluid limits, multiple medications, and regular dialysis sessions. Many patients struggle to adhere consistently. The NANDA diagnosis "Ineffective Therapeutic Regimen Management" describes a pattern of difficulty integrating treatment into daily life.

Defining characteristics: Patients may miss dialysis appointments, fail to follow dietary guidelines, skip medications, or show poor knowledge about their condition. They may express frustration, hopelessness, or a sense of being overwhelmed.

Related factors: Cognitive decline from uremia, depression, lack of social support, financial constraints, and inadequate health literacy all contribute. The sheer complexity of the regimen can be daunting.

Nursing interventions: A nonjudgmental, patient-centered approach is essential. Nurses should assess barriers to adherence and work collaboratively to find solutions. Simplifying medication schedules, providing written instructions, and involving family members can be effective. Motivational interviewing helps patients explore their own reasons for change. Referral to a dietitian, social worker, or counseling services may be appropriate.

DEVELOPING A COMPREHENSIVE CARE PLAN

Using multiple NANDA nursing diagnoses for kidney failure allows the nurse to create a holistic care plan that addresses the patient's medical, emotional, and social needs. For example, a patient with stage 5 chronic kidney disease starting hemodialysis might have the following diagnoses:

- Excess Fluid Volume related to decreased renal function
- Risk for Electrolyte Imbalance related to impaired excretion
- Fatigue related to anemia and uremia
- Risk for Infection related to vascular access device
- Ineffective Therapeutic Regimen Management related to new diagnosis and complex self-care demands

Each diagnosis is accompanied by specific goals and interventions. Goals should be measurable and realistic. For instance, a goal for Excess Fluid Volume might be: "Patient will maintain fluid weight gain of less than 1 kilogram between dialysis sessions within one week." The nurse then implements

interventions such as daily weights, fluid education, and collaboration with the dietitian. Evaluation occurs continuously, with adjustments as needed.

THE ROLE OF PATIENT EDUCATION

Patient education is a thread that runs through every NANDA nursing diagnosis for kidney failure. Without a solid understanding of their condition and treatment, patients cannot participate effectively